

Name: _____

Date/Time: _____

Arrival Time: _____

Date Of Birth: _____

JAMES D. TOROSIS, MD

VICKY W. YANG, MD

DANIEL S. RENGSTORFF, MD

CYNTHIA W. LEUNG, MD

LOCATION: _____

COLONOSCOPY WITH Miralax/Gatorade Prep AM/4 pills Preparation Instructions

Purchase these over-the-counter items at your pharmacy:

- One bottle of **Miralax** powder 238 grams
- Small box of **Dulcolax (bisacodyl)** laxative tablets
- One 64 ounce bottle of **Gatorade Thirst Quencher**. (Avoid red colors)

Five days before the procedure:

Please do not take any coumadin, aspirin, or nonsteroidal anti-inflammatories, such as ibuprofen (Advil, Motrin), naproxen (Aleve), Feldene, Relafen, etc. **Acetaminophen (Tylenol) is acceptable.** ***If you have a history of blood clots, coronary artery stents or a mechanical heart valve please discuss this with your doctor.***

- Please stop eating food products containing seeds and Iron supplements 5 days prior to the procedure.

One day before the procedure:

You may not have any solid foods. You may have clear liquids only. These include: coffee/tea with sweetener only, beef/chicken broth, apple juice, white grape or white cranberry juices, Gatorade, soda pop, popsicles, and Jell-O. Please, no red or purple colors.

At 1PM, take 2 Dulcolax tablets with an 8oz glass of water. Swallow the pills whole. Do not crush or chew. You may experience bowel movements 1-6 hours later.

At 3pm, take 2 additional Dulcolax tablets with 8 oz water.

At 5 PM, mix the 238 gram bottle of Miralax in the 64 oz. bottle of Gatorade. Shake solution until the Miralax is dissolved. Drink an 8 ounce glass about every 15 minutes until completely gone. This will take at least 2 hours. For dinner this evening, you may have any of the above clear liquids.

STOP DRINKING ALL LIQUIDS:

For a **morning** procedure: Do not eat or drink anything after midnight the night before.

You may take your morning medications with a small sip of water, unless otherwise instructed. If you are a diabetic, please discuss your medications with one of us.

You will most likely receive sedation and will require a designated driver to take you home. Public transportation and taxis are **not an option**. If you need assistance, our office may be able to help you find a ride.